



SOCIETY FOR PUBLIC HEALTH EDUCATION

10 G St., NE, Suite 605 • Washington, DC 20002-4242 • Phone 202/408-9804 • Fax 202/408-9815 • Website www.sophe.org

**HEB SELF-STUDY
CONTINUING EDUCATION CREDITS
EVENT 01 HEB 195
Vol. 38 No. 3 June 2011**

Name: _____

Address: _____

Telephone (Home) _____ (Work) _____

E-Mail: _____ CHES# _____

Title: “The Healthy Teen Girls Project: Comparison of Health Education and STD risk reduction intervention for incarcerated adolescent females

Angela R. Robertson, Janet St. Lawrence, David T. Morse, Connie Baird-Thomas, Hui Liew and Kathleen Gresham

1. Existing studies of juvenile offenders offer little evidence of:
 - A. HIV Prevention Knowledge
 - B. Intentions to Use Condoms
 - C. Behavioral Changes
 - D. Condom Attitudes

2. The measures used for self-reported condom use and sexual behavior used in this study have not been used in studies involving which of the following:
 - A. Incarcerated Male Adolescents
 - B. HIV Risk Reduction Interventions with African American Adolescents
 - C. Substance Abusing Youth
 - D. Inner City Latino Adolescents

3. Approximately what percentage of participants reported being a member of a gang?
 - A. 20%
 - B. 24%
 - C. 30%
 - D. 15%

4. What STD(s) were participants screened for at baseline and at follow-up?
 - A. Syphilis & Gonorrhea
 - B. Gonorrhea & Chlamydia

- C. Chlamydia & HIV
 - D. HIV & Herpes
5. Which of these is NOT one of the five outcome variables on intervention effectiveness as developed by St. Lawrence (1993)?
- A. Perceived barriers to condom use
 - B. Sex under the influence
 - C. Alcohol and drug use
 - D. Unprotected sex occasions
6. True or False: The majority of the participants reported marijuana and alcohol use in the six months preceding the trial?
- A. True
 - B. False
7. True or False: There were significant differences between the health education and risk reduction interventions in incidents of STIs post intervention.
- A. True
 - B. False
8. Having an older partner is associated with earlier sexual initiation, less condom use, greater likelihood of STI infection, and pregnancy. What percentage of the participants reported having a boyfriend that was 3 or more years older than them after their release?
- A. 35%
 - B. 38%
 - C. 48%
 - D. 53%
9. Which of the following was NOT given as a possible explanation for the study results?
- A. Cross contamination of the interventions when girls who previously participated reentered the facility during a time of alternative delivery
 - B. There were no differences in condom application and social competency skills between study participants after completing the interventions
 - C. Juvenile offenders have high rates of comorbid mental health and substance abuse problems
 - D. The girls returned to very dysfunctional home after release from incarceration
10. What data analyses techniques did this study use for the statistics section?
- A. Chi-square and ANOVA
 - B. ANOVA and Hierarchical linear modeling
 - C. Linear Regression and Structural Equation Modeling
 - D. Logistic Regression and Chi-square

